



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

OBSERVATIONS *made on the* DISAPPEARANCE *and*
 REAPPEARANCE *of* SATURN'S RING *in the Year*
 1789, *with some Remarks on his diurnal Rotation.* By the Rev.
 H. USSHER, D. D. M. R. I. A. and F. R. S.

THIS year has been remarkably unfavourable in this country to astronomical observations in general, but more particularly so to those proposed to be made on the disappearances and reappearances of Saturn's ring. Read December 5, 1789.

THE first disappearance of the ring in the month of May could not from the state of the weather be observed here at all.

AUGUST 21st. I observed Saturn with the greatest power of the transit instrument; the ring itself was invisible, but its shadow was visible upon the body, though very faint.

AUGUST 26. Observed him with the same power, the ring still invisible.

AUGUST 27. Observed him with a forty-inch double object-glass, before he was near the meridian; ring still invisible, at least with that power, viz. 61 *, which power I ascertained accurately: rain came on before he passed the meridian, and the weather continued rainy and cloudy till the 30th.

AUGUST 30. The ring was visible, particularly on the east side, with the forty-inch achromatic mentioned above: the same also with the greatest power of the transit instrument. By comparing the gradual increase (as far as I can trust to my recollection) I imagine that the ring became visible some time in the night of the 28th; I mean to one using the same instruments that I did.

OCTOBER 1. The ring was just visible on the eastern side only with the double achromatic, and likewise with the transit instrument. Storm came on and continued to the 3d, on which night I got an observation, but the ring was not visible with the transit instrument; there was a heavy gale of wind.

MR.

* It is much to be regretted that the generality of observers do not mention particularly that they have ascertained the power of their telescopes by proper experiments, particularly in observations of this sort. I measured mine by two different methods, viz. by the breadth of the emergent pencil, and also by the usual method of a distant circle and near parallel lines, the latter method gave 58, the other 61; this I believe best.

MR. Newenham of Cork informed me by letter that he saw it on the eastern side to the 5th, when at 10 H. 30' he lost sight of it. This observation he made with a reflecting telescope of seven feet focus, and power 300.

THE power I used was superior to this ; but the transit instrument being fixed, is badly suited to this kind of observation, as the planet passes the field with great rapidity, and with this instrument one cannot pursue it ; which is a circumstance essentially necessary to distinct vision, as I formerly mentioned to this Academy in a paper published in the first volume of our Transactions.

THE observations which I have made are hitherto so badly circumstanced, that any deduction from them with respect to the tables, &c. would be trifling, until the next reappearance, which may either invalidate or fortify such as have been mentioned.

It is, however, worthy of remark at present, that Saturn now divested of his ring appears exceedingly oblate ; much more so than I could have ever supposed from common observation whilst his ring was visible : and as his rotation has not, that I know of, been yet determined by any spots visible upon his surface, it occurred to me that if his equatorial and polar diameters were accurately measured, I could infer his time of rotation by means of the 19th Proposition of Book III. of Newton's Principles, by a deduction from the formula which he there employed, to find
the

the proportion of the polar and equatorial dimeters of Jupiter, by comparison of the density and rotation of the earth with those of Jupiter.

LET T^2 = the square of the time of the fidereal rotation of the earth; and t^2 = the square of that of Saturn, the root of which is sought: let Δ = the earth's density, and δ that of Saturn; D = Saturn's greater diameter, and d his lesser one. Then from Sir Isaac Newton's formula I deduce $t = \sqrt{\frac{d \times \Delta \times 1 \times 1}{\delta \cdot 229 \cdot D - d}} \times 23'.56''$

THE micrometers ordered for the Observatory being not yet arrived, I requested a gentleman of known accuracy to take these measures for me; he was so obliging as to send me a great number, agreeing remarkably together, from which I find Saturn's diameters, reduced to his mean distance, $18''.12$ and $15,855$.

FROM hence, taking Sir Isaac Newton's ratio of the earth's equatorial diameter and axis, and that of the earth and Saturn's densities, as by him computed, the formula will give for Saturn's fidereal rotation 10 H. $12'\frac{1}{2}$.

It is a circumstance worth remarking, that the celebrated Huyghens, in his whimsical and ingenious work, intituled Cosmotheoros, has the following passage: " Quam habeant dierum longitudo (Saturnicolæ sciz.) certo cognosci nequit; sed ex comitis intimi distantia ac periodo, exque eorum comparatione cum intimo Jovialium; verisimile fit non longiores esse dies
" illas

“ illas quam sint in Jove sciz. 10 horarum aut paulo minus.”

We dwell with pleasure upon the lightest conjectures of such an enlightened mind.

By the same formula I computed the time of his rotation, taking the density of Saturn as deduced by Mr. de la Lande in the fourth volume of his *Astronomy*, by which it came out 12 H. 55 $\frac{1}{3}$.

I COMPUTED it also, taking this last density and Mr. Bouguer's ratio of the earth's diameters (which has many advocates) and on this supposition it came out 14 H. 44 $\frac{1}{2}$ nearly.

FROM the great modern improvement of telescopes it may reasonably be expected that his time of rotation will soon be determined by actual observation; and from the two satellites lately discovered by the celebrated Mr. Herschel, his density will soon be more accurately ascertained.

AND when the density of the Georgian planet shall have been determined by means of his satellites, discovered also by Mr. Herschel, perhaps some correct law of the densities may be had; which, if it be found to obtain accurately in those planets which have satellites, may, perhaps reasonably, be extended to those which have none; and enable us to determine the quantity of matter and density of Venus, Mars and Mercury, without having recourse to the decrement of the obliquity of the ecliptic, a matter so delicate in itself, that even at this day,
there

there are some who doubt the fact, though supported by sound theory and observation.

THE following method of determining the quantity of matter in Mars, Venus and Mercury, or any other planet, is certainly true in theory.

THE axis major of the earth's orbit as it revolves round the common center of gravity of earth and sun = 200000. The quantity of matter in the earth we assume = 1, that of the sun is then 352813, the parallax being allowed 8".6. Now by Prop. 60, Book I. of Newton's Principles, as $\sqrt[3]{352814} : \sqrt[3]{352813}$ so is the axis major of the ellipse round the common center of gravity, to the axis major of an ellipse which would in the same periodical time be described round the sun at rest, and is therefore had by the analogy.

VENUS's axis major (Coff. El. Ast.) round the common center of gravity of the sun and Venus = 144662 by observation. Now in the subsequeplicate ratio of the periodical times of the earth and Venus take the earth's diminished axis major to a fourth quantity, and we have the axis major of the ellipse which Venus would describe round a fixed sun; but we had it by observation round the common center of gravity.

THEN let \odot = the sun's quantity of matter, φ = that of Venus, let m = the axis major of her orbit round the common center of gravity, and I = that deduced as above round the fixed sun.

Then

Then $m : I :: \sqrt{\odot + \varphi} : \sqrt{\odot}$. And $m : I :: \odot + \varphi : \odot$. Therefore $m - I : I :: \varphi : \odot$. And therefore $I : m - I :: \odot : \varphi$. Q. E. I. from whence the density is easily deduced. However I freely allow all the difficulties of reducing this to practice.

N. B. If we take the law of the densities laid down by Mr. de la Lande, viz. as the roots of mean motion, the density of Saturn instead of ,10448, which is his number, will come out ,18351, much nearer to Newton's, which reduced to this scale would be ,16750. But Newton's quantities of matter were necessarily defective from applying the solar parallax $10''.5$ instead of $8''.6$.